





**Original Article** 

**Open Access** 

# Presenting a causal model of self-awareness in patients with multiple sclerosis based on visual-spatial development and perceived physical fitness; the mediating role of self-control (application of path analysis)

\*Mehdi Yousefvand<sup>1</sup>; Ahmad Haghi<sup>2</sup>; Zahra Ziar<sup>3</sup>; Zolaikha Zarei<sup>4</sup>

<sup>1</sup>Department of Educational Sciences, Farhangian University, Tehran, Iran.
<sup>2</sup>Faculty member of Psychology Department, Delfan Higher Education Center, Lorestan University, Iran.
<sup>3</sup>M.Sc. in general psychology, Islamic Azad University, Branch of Borujard, Borujard, Iran.
<sup>4</sup>M.Sc. in educational psychology, Payam Noor University of Javanroud, Javanroud, Iran.

## Abstract

**Introduction:** The present study aimed to present a causal model of self-awareness in Multiple Sclerosis (MS) patients based on visual-spatial development and perceived physical fitness with the mediating role of self-control.

**Materials and Methods:** The statistical population of this correlational study included all patients with MS in Borujerd, Iran, in 2023-2024, of which 300 people were selected by convenience sampling. The questionnaires of visual-spatial development, perceived physical fitness, self-control, and self-awareness were used to collect data. The data were analyzed using SPSS 24 and AMOS software.

**Results:** The findings showed that visual-spatial development affected self-control ( $\beta$ = 0.54 and value 8.09) and self-awareness ( $\beta$ = 0.67 and value 11.76); perceived physical fitness had an effect on self-control ( $\beta$ =0.59 and value 9.96) and self-awareness ( $\beta$ = 0.55 and value 8.85) and self-control affected self-awareness ( $\beta$ = 0.47 and value 6.70) (P< 0.01). The findings also showed that the present model explained 33 percent of the variance in self-control and 42 percent of the variance in self-awareness. In addition, self-control plays a role as a mediating variable in the relationship between exogenous variables and self-awareness.

**Conclusion:** According to the results, increasing self-control and consequently increasing the level of visual-spatial development and perceived physical fitness will improve the level of self-awareness in patients with MS. This suggests that relevant authorities should pay more attention to the role of self-control as a mediating variable.

Keywords: Perceived physical fitness, Self-awareness, Self-control, Visual-spatial development

#### Please cite this paper as:

Yousefvand M, Haghi A, Ziar Z, Zarei Z. Presenting a causal model of self-awareness in patients with multiple sclerosis based on visual-spatial development and perceived physical fitness; the mediating role of self-control (application of path analysis). Journal of Fundamentals of Mental Health 2025 Mar-Apr; 27(2):79-86. DOI:10.22038/JFMH.2025.82054.3159

\*Corresponding Author:

Department of Educational Sciences, Farhangian University, Tehran, Iran. mehdi\_yousefvand@cfu.ac.ir Received: August. 23, 2024 Accepted: Jan. 12, 2025

Copyright © 2025 Mashhad University of Medical Sciences. This work is licensed under a Creative Commons Attribution-Noncommercial 4.0 International License https://creativecommons.org/licenses/by-nc/4.0/deed.en

Multiple sclerosis (MS) is an inflammatory disease in which the myelin sheaths of brain and spinal cord nerve cells are damaged (1). MS is a disease related to the central nervous system (2,3). MS is most common in young people aged 20 to 40 years and is three times more common in women than in men (4,5). The most common symptoms of MS are fatigue, dizziness, loss of balance or numbness, blurred vision, slurred speech, prolonged double vision, sexual problems, and sensitivity to cold (6.7). According to the MS Association, concern about the future is particularly important for patients (8). Due to its unpredictable nature 9), this disease can disrupt the quality of life of the affected person (10), making it difficult to carry out daily activities (11).

Studies have shown that the level of selfawareness in MS patients undergoes many changes (12). The better patients know themselves, the more likely they are to overcome anxiety (13,14) and live a normal life (15-17). In other words, self-awareness in MS patients provides the basis for self-efficacy to control negative emotions (18,19). According to neuropsychological theorists, self-awareness is the nervous system's ultimate understanding of its nature (20-22). In psychology, selfawareness means understanding all thoughts, beliefs, strengths, and weaknesses (23,24) and helping to improve oneself in challenging situations (25). Self-awareness in MS patients has also been defined as a developed sense of mindfulness (26), perceived physical health (27), and basic self-esteem (28).

Researchers have shown that physical fitness exercises affect the level of self-awareness and mindfulness (29,30) and psychological health of MS patients (31,32). Central neuroscience researchers have shown that visual-spatial processing in MS patients undergoes many negative changes (33). People with MS are involved in psychosomatic illness and have problems in the field of visual-spatial perception and physical fitness (34). Visualspatial processing is one of the most important processing foundations and forms mental representations about the outside world (35). Visual-spatial processing includes all spatial information, which is unfortunately severely impaired in MS patients (35). Research has shown that MS patients have problems in processing motion perception, shape and background perception, depth perception, and

visual-motor coordination (35). Some studies have shown that MS patients have poor perceived physical fitness and self-assessment (36). The perception of MS patients' physical fitness directly affects their self-awareness (37). It is important to note that perceived physical fitness can help MS patients to have high self-control and self-awareness (38) by improving physical imagery (39).

Psychologists also believe that perceived physical fitness in MS patients helps to control and overcome the disease (39). Therefore, it is clear that MS patients who increase their level of perceived physical fitness will have a more positive attitude, higher self-awareness, and more self-control in MS and will be able to cope with this disease better and more favorably (32,33). Other studies have reached interesting and, of course, different results. One of these findings is that the self-control construct (16,18,21,27,34,36,38) mediates the relationship between influential and predictive variables of self-awareness. such as visuospatial development and perceived physical fitness in MS patients. Researchers have recently investigated the effect of selfimplementing mindfulness control in techniques to accelerate the psychological wellbeing of MS patients (39). Therefore, and in light of the above, the present study aims to present a model of self-awareness based on visuospatial development and perceived physical fitness and the mediating role of selfcontrol in MS patients.

### Materials and Methods

The present research design was based on the correlational approach using path analysis. The statistical population included all women with MS in Borujerd, Iran, in 2023-2024 (n= 1561) who were selected using the convenience sampling method. For studies based on model drawing, the following rule of thumb has been suggested for sample selection: The minimum sample size for each calculated parameter is 5, a ratio of 10 to 1 is considered more appropriate, and a ratio of 15 to 1 is considered desirable. Since there are 20 observed parameters in the present study, a ratio of approximately 15 to 1 was used, and 300 people were selected using the convenience method. The inclusion criteria were age over 35, no debilitating diseases or chronic heart and respiratory diseases, no participation in treatment programs, and being literate and willing to participate in the research.

The exclusion criteria were lack of participation and cooperation in the research implementation process.

## Research instruments

A) Visual-Spatial Development Questionnaire: This questionnaire, developed by Harry and Wider, has 51 questions and six subscales: body awareness and sense, body location in space, body-environment relationship and environment-other relationship. spatial maintenance, visual, logical reasoning, and representational thinking. The three-point scoring range is from never=3 to always=1. The minimum and maximum scores in the entire questionnaire are 51 and 153, respectively. This questionnaire has been assessed as appropriate in the content, face, and criterion validity project (40).

*B)* Perceived Physical Fitness Questionnaire: The questions of this tool are scored on a 5-point Likert scale from strongly disagree= 1 to strongly agree= 5. This questionnaire has 12 questions and four sections: physical condition (questions 1 to 3), flexibility (questions 4 to 6), muscle condition (questions 7 to 9), and body composition (questions 10 to 12). The minimum and maximum scores in the entire questionnaire are 12 and 60, respectively. Cronbach's alpha for the entire instrument is reported to be 0.88 and is desirable (41).

C) Self-awareness questionnaire: This scale was developed by Cooper and consisted of 25 questions. The questionnaire has three subscales: private self-awareness (questions 1 to 10), general self-awareness (questions 11 to 18), and social self-awareness (questions 19 to 25), which are set on a 5-point Likert scale from never= 1 to very much= 5. The minimum score in this instrument is 25, and the maximum is 125. Cronbach's alpha for the entire instrument was reported to be 0.75 and desirable (42).

After obtaining the necessary permits, the researchers referred to a comprehensive health center in Borujerd dedicated to providing services to MS patients. A list of all patients was provided to the researchers, who were selected from a list of 300 people using the available method. In the next stage, they were contacted based on their psychiatric and medical records, including their ID card details, address, and home phone number.

After contacting them, the process of providing the questionnaires began, along with the study objectives and obtaining consent. Filling out the questionnaires was done in two ways. In the first method, women with MS answered the questionnaires in person. In the second method, some MS women who could not come to the center in person were sent the questionnaires online, and they answered them. To be precise, 293 participants answered the questionnaires accurately and completely. In other words, seven people did not answer the questions completely, and these seven people were excluded from the analysis.

The present study used descriptive statistical methods to analyze the collected data. SPSS24 and AMOS24 software were used for quantitative statistical analyses.

# Results

Demographic findings showed that 53% of the participants were between the ages of 20 and 30, and 47% were between the ages of 31 and 50. In addition, findings related to employment status, family economic status, and education level showed that 71.66% of women were employed and 28.33% were unemployed.

Also, 38.33% of MS women had low economic status, 40.00% had medium economic status, and 21.66% had high economic status. In addition, 26% of MS patients had a diploma or lower diploma education, 50.66% had a postgraduate diploma or bachelor's degree, and 23.33% had a postgraduate degree or higher.

Table 1 reports descriptive findings related to the mean and standard deviation of the variables. Based on the Pearson correlation coefficient method, it was determined that the self-awareness variable has a positive and significant relationship with visual-spatial development (P < 0.05, r= 0.57) and mental balance (P < 0.05, r = 0.44). In addition, the correlation between visual-spatial development and mental balance is (P < 0.05, r = 0.42). The results showed that the assumptions of normality, multivariate normality (Mardia coefficient= 4.78), multiple non-collinearity, univariate outliers (using boxplots), and multivariate outliers (using Mahalanobis distance) were met. The results of the Bootstrap method showed that the path of visual-spatial transformation to self-awareness mediated by mental balance was in the range of the lower limit= 0.215 and the upper limit -0.302 at a significance level of 0.005. In this study, the assumption of linearity was confirmed using the scatterplot method.

Table1. Descriptive indicators in the variables of visual-spatial development, mental balance, and self-

awareness									
Variable	Subscale	Mean	Standard Error	Skewness		Skewness		Index of Multiple Collinearity	
				Index	Standard Deviation	Index	Standard Deviation	Tolerance Factor	Inflation Factor
Visual- spatial developme nt	Representation al thinking	21.45	2.44	1.74	0.14	2.43	0.25	0.47	2.22
	Visual logical reasoning	19.80	2.40	1.08	0.14	2.22	0.25	0.70	1.75
	Maintaining space Body	20.49	2.89	2.72	0.14	0.74	0.25	0.66	1.10
	awareness and	21.82	2.71	0.99	0.14	0.53	0.25	0.49	2.20
	Location of the body in space Relationship of	20.21	2.90	0.99	0.14	0.53	0.25	0.72	1.11
	the environment with others and body with the environment	22.73	2.14	0.52	0.14	0.20	0.25	0.70	1.36
	Visual-spatial development (total score)	121.67	11.83	1.03	0.14	0.56	0.25	0.74	1.83
Mental balance	Experiential acceptance	29.30	4.21	3.09	0.11	11.09	0.23	0.42	2.29
	Non-reaction Mental balance	36.48 100.96	5.14 9.12	1.41 1.23	0.11 0.11	1.40 0.73	0.23 0.23	0.73 0.74	1.71 1.75
Self- awarenes	Private self- awareness Public self- awareness	33.55	3.25	1.87	0.15	0.70	0.26	0.70	1.11
		36.32	4.10	1.80	0.15	0.79	0.26	0.63	1.29
	Social self- awareness Self-awareness	22.87 96.15	3.85 10.28	1.81 1.86	0.15	0.77 0.74	0.26	0.50	2.32
	Sen-awareness	70.15	10.20	1.00	0.15	0.7	0.20	0.07	1.10



Figure 1. Model in standard coefficients mode

As shown in Figure 1, the present model explains 33% of the variance in self-control, 30% in mental balance, and 42% in self-awareness. The results showed that visual-spatial development had an effect on self-control with an impact coefficient of 0.54 and a value of 8.09; on mental balance with an impact coefficient of 0.60 and a value of 9.58; and on self-awareness with an impact coefficient of 0.67 and a critical value of 11.76. The results also showed that perceived physical fitness had an effect on self-control, with an impact coefficient of 0.59 and a value of 9.96; on mental balance, with an impact coefficient of 0.46 and a value of 6.58; and on self-awareness,

with an impact coefficient of 0.55 and a critical value of 8.85. In addition, the results showed that sports mindfulness affected self-control, with an impact coefficient of 0.50 and a value of 7.10; mental balance, with an impact coefficient of 0.59 and a value of 9.97; and self-awareness, with an impact coefficient of 0.51 and a critical value of 64.7; self-control had an impact on self-awareness with an impact coefficient of 0.47 and a value of 70.6, and mental balance had an impact on self-awareness with an impact on s

Table 2. Parameters	for measuring	direct relationships	between variables	in the proposed mod	el and model fit
		• •	•		

indices								
Path	SD	В	β	CR	α			
visual-spatial development on mental balance 0.60		0.65	0.60	3.47	0.001			
visual-spatial development on self-awarenes	spatial development 0.67 0.70 n self-awarenes		0.67	1.75	0.001			
mental balance on self- awarenes	0.50	0.63	0.57	1.24	0.001			
Model fit indices								
indices		F		Accepted domain	Result			
$(X^2)$		426.42		> 0.05				
(P.value)		0.001			Non Desirable			
$(X^2/DF)$		2.76		< 5	Desirable			
(GFI)	0.85			> 0.90	Desirable			
(RMSEA)		0.042		< 0.1	Desirable			
(CFI)		0.91		> 0.90	Desirable			
(TLI)		0.90		> 0.90	Desirable			
(NFI)		0.87		> 0.90	Desirable			
(PNFI)		0.74		> 0.50	Desirable			

#### Discussion

The present study aimed to test the fit of the causal model of self-awareness in MS patients based on visuospatial development and perceived physical fitness with the mediating role of self-control. The results showed that the direct effect of visuospatial development on self-control and self-awareness is significant. This finding is consistent with the results of studies in terms of results (7,11,22). The self-control construct in Clinton et al.'s theory is generally understood as the ability to change or adapt oneself to the world around one's body to

create a better fit between oneself and the world. On the other hand, the concept of visuospatial development has an important subcomponent called the relationship between the individual and the environment.

Therefore, one of the important reasons for explaining why the direct effect of visuospatial development on self-control was significant is the existence of a common element of trying to connect the individual with the environment optimally (13). Regarding the effect of visualspatial development on self-awareness, one of the important explanations is that there is an element of self-attention and review of mental, intellectual, and physical conditions in visualspatial development and self-awareness. Another explanation, in agreement with the studies of Carletto (7), Dietmaier (11), and Kang (22). Our findings showed that the direct effect of perceived physical fitness on selfawareness and self-control is significant. This finding is consistent with the results of studies. Studies have shown that people who are more physically fit also have a higher level of selfawareness. This is because people with higher physical fitness care more about their physical and mental health, and while in the concept of self-awareness, attention to the health of the individual's personality and identity is also important. Therefore, in general, attention to the health of the body and mind in both the concept of perceived fitness and the concept of self-awareness is the main reason that perceived physical fitness has been able to have a causal and direct effect on the concept of selfawareness. One of the main reasons that perceived physical fitness has been able to significantly predict self-control is that in the structure of physical fitness, there are concepts such as attention to physical health, sports management, and regular exercise. In contrast, in the concept of self-control, attention to physical health and daily life is of particular importance (10,18,30,33).

Another explanation, consistent with the studies of Dickstein (10), Hobson (18), Mertens (30), and Ogunwale (33) which conducted on women. In addition, the sample size in the studies of Mertens (30) and Ogunwale (33) was 300, similar to the present study.

The results showed that self-control significantly mediates the relationship between visuospatial development and perceived physical fitness with self-awareness. This finding is consistent with the results of studies (16,27,34,38). In the self-awareness theory of MacRae, it is believed that people who are involved in MS should have a higher level of self-awareness about their disease so that they can better cope with control and follow up on treatment. On the other hand, people who have higher perceived physical fitness and better visual-spatial development have higher selfawareness because high perceived physical fitness and higher self-control and visualspatial development make people control their emotions better and have more optimal arousal levels. So, self-control can mediate the relationship between perceived physical fitness and visual-spatial development with selfawareness and indirectly and significantly affect self-awareness. So, in general, and based on theoretical and empirical foundations, it can be said that the existence of common elements in the concepts of perceived physical fitness, visual-spatial development, and self-control has caused self-control to have a mediating and positive role in the relationship between perceived physical fitness and visual-spatial development with self-awareness (33). Another explanation, in agreement with the studies of Han (16), London (27), Potoczny (34), and Shi (38). One limitation of the present study is that it was conducted on women with MS, and the results cannot be generalized to male patients. Given the mediating role of self-control in the relationship between visuospatial development and perceived physical fitness with selfawareness, it is suggested that the importance of this concept be further emphasized.

## Conclusion

Based on the findings, self-control mediates the relationship between visual-spatial development, perceived physical fitness, and self-awareness, significantly. Therefore, paying attention to these variables is crucial in increasing the level of self-awareness of MS patients.

### Acknowledgments

We hereby express our gratitude to those who helped us conduct this research.

### **Conflict of Interest**

The authors declare no conflict of interest.

#### Funding

No funding

### **Ethical Considerations**

This article is based on the master's thesis at the Islamic Azad University, Boroujerd Branch.

#### **Code of Ethics**

IR.MEDAPAU.REC.1398.015

### **Authors' Contributions**

The design and implementation were carried out by Mehdi Yousefvand and Ahmad Haqi; Mehdi Yousefvand carried out the introduction and statement of the problem; Ahmad Haqi carried out the methodology, discussion, and conclusion; and Zahra Ziyar and Zolaikha Zarei carried out the findings and general review of the work under the supervision of Mehdi Yousefvand.

#### References

1. Ambretti A, Fogliata A, Di Palma D. Innovation in physical education: Proprioception, periferical vision, self-awareness, and sustainability. J Hum Sport Exerc 2024; 19(3): 915-29.

2. Kahl PT. Domain-general abilities and mathematical achievement: A developmental perspective on the roles of executive functions, visual-spatial skills, and emotion regulation. Ph.D. Dissertation. Switzerland: University of Basel, 2021.

3. Bademli K, Lök N. Relationship between the health perception and physical activity of individuals. Turkish journal of sport and exercise 2019; 19(3): 915-29.

4. Soldatova GU, Vishneva AE, Koshevaya AG. [Characteristics of the cognitive domain of children and adolescents with different levels of media tasks]. Russian psychology 2022; 1(3): 36-37. (Russian)

5. Brytek-Matera A, Kozieł A. The body self-awareness among women practicing fitness: A preliminary study. Polish psychological bulletin 2015; 46(1): 104-11.

6. Carden J, Jones RJ, Passmore J. Defining self-awareness in the context of adult development: A systematic literature review. J Manag Educ 2022; 46(1): 140-77.

7. Carletto S, Cavalera C, Sadowski I, Rovaris M, Borghi M, Khoury B, et al. Mindfulness-based interventions for the improvement of well-being in people with multiple sclerosis: A systematic review and metaanalysis. Psychosom Med 2020; 82(6): 600-13.

8. Chen S, Wei H. Linking temporal landmarks to voluntary simplicity: The mediating roles of self-transcendence and self-enhancement. J Bus Ethics 2023; 188(4): 693-708.

9. Davis BE, Lakin L, Binns CC, Currie KM, Rensel MR. Patient and provider insights into the impact of multiple sclerosis on mental health: A narrative review. Neurol Ther 2021; 34(10): 99-119.

10. Dickstein E. Self and self-esteem: Theoretical foundations and their implications for research. Hum Dev 1977; 20(3): 129-40.

11. DietmaierJM, von dem Knesebeck O, Heesen C, Kofahl C. Personality and its association with self-management in multiple sclerosis. Mult Scler Relat Disord 2022; 12(5): 61-76.

12. Gomelsky A, Stolzle AC, Armstrong WP. Pathophysiology of female micturition disorders. In: Female genitourinary and pelvic floor reconstruction. Cham: Springer International Publishing: 71-84.

13. Gontkovsky ST. Understanding visual-spatial perceptual deficits in individuals with multiple sclerosis: An analysis of patient performance on the Hooper Visual Organization Test and visual form discrimination. Int J Neurosci 2023; 133(6): 636-41.

14. Grippa E, Sellitto M, Scarpazza C, Mattioli F, di Pellegrino G. Multiple sclerosis reduces sensitivity to immediate reward during decision-making. Behav Neurosci 2017; 131(4): 325-42.

15. Hamaideh SH, Abuhammad S, Khait AA, Al-Modallal H, Hamdan-Mansour AM, Masa'deh R, et al. Levels and predictors of empathy, self-awareness, and perceived stress among nursing students: A cross sectional study. BMC Nurs 2024; 23(1): 131-9.

16. Han A. Mindfulness-and acceptance-based interventions for symptom reduction in individuals with multiple sclerosis: A systematic review and meta-analysis. Arch Phys Med Rehabil 2021; 102(10): 2022-31.

17. Herson M, Kulkarni J. Hormonal agents for the treatment of depression associated with the menopause. Drugs Aging 2022; 39(8): 607-18.

18. Hobson RP, Chidambi G, Lee A, Meyer J, Müller U, Carpendale JIM, et al. Foundations for self-awareness: An exploration through autism. Monogr Soc Res Child Dev 2006; 12(89): 36-49.

19. Jakimovski D, Benedict RH, Weinstock-Guttman B, Ozel O, Fuchs TA, Lincoff N, et al. Visual deficits and cognitive assessment of multiple sclerosis: confounder, correlate, or both? J Neurol 2021; 268(7): 2578-88.

20. Jhang JF, Jiang YH, Kuo HC. Discriminating different bladder and bladder outlet dysfunctions by urinary biomarkers in women with frequency-urgency syndrome. Biomedicines 2023; 11(3): 673.

21. Jo YG, Kim JS, Yang CH. The effect of self-management and exercise immersion on physical self-awareness among the elderly participating in sports Activities. Ilkogretim online 2021; 20(3): 346-59.

22. Kang W. Understanding the effect of multiple sclerosis on general and dimensions of mental health. J Clin Med 2022; 11(24): 74-83.

23. Lakin L, Davis BE, Binns CC, Currie KM, Rensel MR. Comprehensive approach to management of multiple sclerosis: addressing invisible symptoms—a narrative review. Neurol Ther 2021; 10(8): 75-98.

24. Li C, Yu Q, Wu Y, Fan L, He J. Internet gaming disorder and cognitive failures in daily life among university students: The mediating role of sleep quality and the moderating role of mindfulness. Curr Psychol 2024; 43(38): 30446-58.

25. Liang P, Jiang H, Wang H, Tang J. Mindfulness and impulsive behavior: Exploring the mediating roles of self-reflection and coping effectiveness among high-level athletes in Central China. Front Psychol 2024; 15(56): 130-41.

26. Liao YC, Huang TY, Lin SH, Wu CH, Chang KT, Hsieh S, Yang CT. Mediating role of resilience in the relationships of physical activity and mindful self-awareness with peace of mind among college students. Sci Rep 2023; 13(1): 10386.

27. London M, Sessa VI, Shelley LA. Developing self-awareness: Learning processes for self-and interpersonal growth. Annual review of organizational psychology and organizational behavior 2023; 10(1): 261-88.

28. Mayo CD. Miksche K, Attwell-Pope K, Gawryluk JR. The relationship between physical activity and symptoms of fatigue, mood, and perceived cognitive impairment in adults with multiple sclerosis. J Clin Exp Neuropsychol 2019; 41(7): 715-22.

29. Mazancieux A, Souchay C, Casez O, Moulin CJ. Metacognition and self-awareness in multiple sclerosis. Cortex 2019; 111: 238-55.

30. Mertens J. Walking to beats in music and metronomes in persons with progressive MS: Preliminary pilot study results. MS. Diassertatin. 2023. Available from: http://hdl.handle.net/1942/41037

31. Morin A. Possible links between self-awareness and inner speech theoretical background, underlying mechanisms, and empirical evidence. J Conscious Stud 2005; 12(4-5): 115-34.

32. Naisby J, Wilson-Menzfeld G, Baker K, Morris R, Robinson J, Barry G. Yoga and multiple sclerosis: Maintaining engagement in physical activity. PLoS ONE 2023; 18(7): 124-50.

33. Ogunwale OR. Mathematics education: Reducing effects of dyscalculia in children and adults with hearing impairment. Rethinking the teaching and learning of mathematics in the pandemic era 2023; 23(1): 131-9.

34. Potoczny W, Herzog-Krzywoszanska R, Krzywoszanski L. Self-Control and emotion regulation mediate the impact of karate training on satisfaction with life. Front Behav Neurosci 2022; 15(45): 802-37.

35. Rai SB. Evolutionary association between self-awareness and self-control. In: Towards inclusive societies. India: Routledge India; 2024: 89-100.

36. Scott Ridley D. Reflective self-awareness: A basic motivational process. J Exp Educ 1991; 60(1): 31-48.

37. Setyaningrum F, Siswantari H. Visual-spatial and intrapersonal intelligence in learning of rampak kendang art for elementary school students. International journal of creative and arts studies 2020; 7(1): 49-60.

38. Shi Y, Qu S. The effect of cognitive ability on academic achievement: The mediating role of self-discipline and the moderating role of planning. Front Psychol 2022; 13(289): 101-23.

39. Simpson R, Simpson S, Ramparsad N, Lawrence M, Booth J, Mercer SW. Effects of mindfulness-based interventions on physical symptoms in people with multiple sclerosis–a systematic review and meta-analysis. Mult Scler Relat Disord 2022; 28(5): 346-50.

40. Harry W, Wider H. Visual-Spatial Development Questionnaire. Toxicol Lett 2014; 230(2): 271-81. 41. Sprung JM, Rogers A. Work-life balance as a predictor of college student anxiety and depression. J Am Coll Health 2021; 69(7): 775-82.

42. Nair BP, Prasad T, Nair SK. Self-awareness trigger leading to authentic leadership: conceptualization and development of reliable and valid self-awareness trigger scale. Dynamic relationship management journal 2021; 10(1): 23-44.