



Original Article

Attention deficit hyperactivity disorder in patients with congenital heart disease

Hassan Mottaghi Moghaddam Shahri¹; Mahboubeh Eslamzadeh²;
*Aazam Sadat Heydari Yazdi³

¹Associate professor of pediatric and congenital cardiology, Pediatric Department, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

²Assistant professor of psychiatry, Psychiatry and Behavioral Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

³Assistant professor of child and adolescent psychiatry, Psychiatry and Behavioral Sciences Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

Introduction: Neurodevelopmental disabilities, particularly executive function impairments, are currently the most prevalent long-term morbidity in the population with Congenital Heart Disease (CHD). This study aimed to investigate the frequency of Attention Deficit Hyperactivity Disorder (ADHD) in children with congenital heart disease.

Materials and Methods: This was a retrospective cohort study, which was performed during 2002-2018 on patients with CHD referred to the pediatric cardiology clinic of Imam Reza hospital, Mashhad, Iran. Using the census method, all files for which ADHD diagnosis should be included in the study. Diagnosis of CHD was performed clinically and using an echocardiographic machine by an experienced pediatric cardiologist. Demographic, clinical, and para-clinical data of patients and the type of heart intervention were collected and analyzed. The data were analyzed through SPSS software version 16.5, descriptive statistics, and the Chi-square test.

Results: In this study, 136 patients were enrolled. The mean age of participants in the study was 59.12± 45.84 months at the time of diagnosis. An abnormal electroencephalogram (EEG) was reported in three cases (2.2%). There was a significant correlation between prematurity and developmental delay ($P= 0.01$). The mean age was significantly different in patients with minor rather than those with major cardiac disorders ($P< 0.05$). The prevalence of ADHD in the CHD population was high (31.6%).

Conclusion: This study demonstrated that children born with congenital heart disease are at increased risk of suffering from attention deficit hyperactivity disorder.

Keywords: Attention deficit hyperactivity disorder, Congenital heart disease, Neurocognitive disorder, Pediatric.

Please cite this paper as:

Mottaghi Moghaddam Shahri H, Eslamzadeh M, Heydari Yazdi AS. Attention deficit hyperactivity disorder in patients with congenital heart disease. *Journal of Fundamentals of Mental Health* 2023 Jan -Feb; 25(1):11-18.

Introduction

Congenital Heart Disease (CHD) is the most common congenital disability worldwide, affecting millions of newborns yearly (1).

The prevalence of CHD between 1970 and 2017 globally was 8.22 per 1000. During this period, the overall prevalence of CHD globally increased by 10% every five years (1). In order

*Corresponding Author:

Psychiatry and Behavioral Sciences Research Center, Ibn-e-Sina Hospital, Mashhad, Iran.

heydarias@mums.ac.ir

Received: Aug. 21, 2022

Accepted: Nov. 22, 2022

to improve CHD management, most patients can be survived to higher age in childhood or even more. There are now challenges related to these patients' quality of life and long-term development. It is established that children with CHD are more susceptible to suffering deficiencies in intellectual functioning, developmental problems, and academic performance difficulties (2,3). Children with CHD are at higher risk of several cognitive disorders, which can show problems in school, like patients with attention disorders (4). Unfavorable effects of chronic and intermittent hypoxia on development, behavior, and academic achievement have been reported in previous studies in children with CHD (5).

Neurodevelopmental disabilities, particularly executive function impairments, are currently the most prevalent long-term morbidity in the population with CHD (6). Executive function refers to a set of higher-order neurocognitive abilities that coordinate and organize actions toward a goal, allowing the individual to adapt to new or complex situations (7). Impairments in executive function manifest as behavioral dysregulation, attention problems, impaired working memory (like the ability to keep information in mind and manipulate it over a short period), and problems with organization and planning abilities. Executive function is more strongly associated with school readiness than IQ, predicts mathematics and reading competence throughout the school years (8,9), and is strongly associated with social cognition (8). Patients with CHD display deficits in visual-perceptive skills (10-14) and executive function (10,11), ADHD symptoms (14,15), and reduced quality of life (16,17). There are rarely available data on mental health outcomes in CHD survivors in adolescence (18,19).

According to full DSM-5 criteria, the prevalence of ADHD was 3.55% in a large-scale study (20). Although long-term behavioral outcomes have been studied for various forms of CHD, the presence of ADHD within this group of patients is rarely addressed (21). Studies conducted with heterogeneous CHD populations have reported increased risk and under-treatment of psychiatric symptoms, including anxiety and depression (19,22). However, the prevalence of psychiatric disorders in adolescents with critical CHD remains under-investigated. Although studies suggest that adolescents with critical CHD display a higher incidence of ADHD (11,15,21),

this literature is limited by reliance on parent- and self-report measures. This study aimed to investigate the prevalence of ADHD in children with CHD referred to Imam Reza hospital, Mashhad- Iran between 2002 and 2018.

Materials and Methods

This was a retrospective cohort study performed during 2002-2018 in all patients with CHD referred to the pediatric cardiology clinic of Imam Reza hospital, Mashhad, Iran.

The inclusion criteria included children and adolescents aged less than 18 years who were diagnosed with CHD (ICD-9-CM codes: 745, 746, 747.1-747.4) by a pediatrician, cardiologist, or cardiac surgeon.

The exclusion criteria included patients with histories of psychiatric disorders (ICD-9-CM codes: 290-319), confirmed diagnosis of an autism spectrum disorder that would prevent successful completion of the planned study testing, and scheduled to undergo major cardiac interventions in the six months following enrollment.

CHD definition

All patients were diagnosed with pediatric and congenital cardiologists according to ICD-10 and met one of the following criteria:

a) Age less than 18 years at baseline assessment, b) had received cardiovascular care at the Imam Reza Hospital, Mashhad, and c) Diagnosis of CHD was performed clinically and using an echocardiographic machine and other cardiac imaging modalities by an experienced pediatric cardiologist. Heart diseases were divided into two important (major) and non-important (minor) categories based on the need for follow-up and intervention.

Echocardiographic assessment

All patients underwent M-mode, two-dimensional, Doppler, and color Doppler echocardiography performed on (Vivid 7 GE, USA, and Resona7, Mindray, China) using 3-5 MHz transducer by an experienced pediatric echocardiographer in the department of pediatric and congenital cardiology of Imam Reza hospital. Structural defects and cardiac function were determined by using standard and oblique views. Attention-deficit / hyperactivity disorder assessment.

The diagnoses are based upon assessments by a psychiatrist or child and adolescent psychiatrist

according to DSM-IV or DSM-V criteria. The institutional review board approved this study at Mashhad University of Medical Sciences, Mashhad, Iran. All parents or guardians were requested to complete the written consent form for enrolling in the study. All data were entered in SPSS software Ver. 16.5.

Descriptive data were reported as count, percent, mean, and standard deviation. For comparing qualitative variables, the Chi-square test was used.

P value less than 0.05 was considered significant.

Results

During the period of study, 136 patients enrolled in the study. The mean age of the participants in the study was 59.12 ± 45.84 months at the time of diagnosis. Twenty-nine were female (21.3%), and others were male (78.7%). Table 1 shows the demographic data of patients.

Table 1. Demographic data of participants

Characteristics	
Age at diagnosis of CHD (months, SD)	59.12 ± 45.84
Sex (n, %)	
Male	107, 78.7
Female	29, 21.3
Age group (n, %)	
< 36 months	44, 32.4
37-80 months	44, 32.4
81-125 months	37, 27.2
126-170 months	9, 6.6
>171 months	2, 1.5
Age at last visit (months, SD)	105.63± 50.45
Birth weight (mean, SD) (gram)	3075± 603.51
Birth weight group (n, %)	
Small for gestational age	24, 17.6
Normal	104, 76.4
Large for gestational age	8, 5.9
New Weight (mean, SD) kilogram	31.25± 15.78
Premature (n, %)	5, 3.7
Consanguinity (n, %)	48, 35.3
Delivery (n, %)	Cesarean (74, 54.4) Vaginal (62, 44.1)
Syndrome (n, %)	Down (3, 2.2), Noonan (2, 1.5)

Table 2 shows referring causes of patients to the pediatric cardiology department, and Table

3 shows the clinical features of patients enrolled in this study.

Table 2. Referring causes of patients to the pediatric cardiology department

Variable	
Referring groups (n, %)	Pediatrics (77, 56.6) Psychiatrists (3, 2.2) Patients (28, 20.6) Others (28, 20.6)
Cause of Referring (n, %)	Clinical suspicion (8, 5.9) Respiratory symptoms (47, 34.6) Abnormal Chest radiography (2, 1.5) ADHD (3, 2.2) Cardiovascular manifestation (75, 55.1) Other (1, 0.7)

Table 3. Clinical features of patients

Characteristics	
Tachycardia (n, %)	4, 2.9
Click or murmur (n, %)	120, 88.2
Anemia	2, 1.5
Sweating	4, 2.9
Chest anomaly	5, 3.7
Icterus	1, 0.7
Hematologic abnormalities	4, 2.9
Gastrointestinal abnormalities	12, 8.8
Neurologic abnormalities	24, 17.6
Ear nose throat abnormalities	4, 2.9
Respiratory abnormality	15, 11
Ophthalmologic abnormalities	3, 2.2
Urogenital abnormalities	8, 5.9
Orthopedic abnormalities	2, 1.5

Two patients had frequent urinary tract infection (1.5%). Cardiac abnormality is listed in

Table 4. Electrocardiogram (ECG) abnormality was reported in 12 cases in Table 5.

Table 4. Major and minor cardiac abnormalities of patients

Major abnormalities	N, %
Septal defects	
Ventricular septal defect	27, 19.9
Atrial septal defect	4, 2.9
Atrioventricular septal defect	2, 1.5
Arterial abnormality	
Patent ductus arteriosus	8, 5.9
Right sided obstructive lesions	
Tetralogy of Fallot	6, 4.4
Pulmonary valve stenosis	5, 7.1
Pulmonary atresia (n, %)	1, 0.7
Pulmonary vessel abnormality	
Pulmonary hypertension	2, 1.5
Left sided obstructive lesions	
Aortic stenosis	7, 5.1
Coarctation of aorta	4, 2.9
Myocardial abnormality	
Hypertrophic cardiomyopathy (n, %)	1, 0.7
Complex	
Double Outlet Right Ventricle (DORV) (n, %)	2, 1.5
Transposition of great arteries (n, %)	1, 0.7
Heterotaxia (n, %)	2, 1.5
Tricuspid valve abnormality	
Ebstein anomaly (n, %0)	1, 0.7
Minor abnormalities	N, %
Floppy mitral valve and/ or mitral valve prolapse without mitral regurgitation	56, 41.1
Patent foramen ovale	12, 8.8
Persistent left superior vena cava	4, 2.9

Table 5. Electrocardiogram abnormalities

Characteristics	
QRS wideness > 0.16 Sec	1, 0.7
Right Bundle Branch Block (RBBB)	5, 3.7
Premature Atrial Contraction (PAC), Premature Ventricular Contraction (PVC) (Infrequent)	1, 0.7
Sinus arrhythmia	3, 2.2
Right Axis Deviation (RAD) and Right Ventricular Hypertrophy (RVH)	2, 1.5
Left Bundle Branch Block (LBBB) and ST-T changes	1, 0.7
Tachycardia	1, 0.7
Right Atrial Hypertrophy (RAH)	1, 0.7

Abnormal electroencephalogram (EEG) was reported in three cases (2.2%). Neuropsychiatric

disorders were reported in 34 cases (25%). Table 6 shows the prevalence of these disorders.

Table 6. Neuropsychiatric disorders of patients

Characteristics	
Bruxism	1, 0.7
Learning disorders	3, 2.2
Autism	1, 0.7
Anxiety	3, 2.2
Enuresis	1, 0.7
Development delay	15, 11
Behavioral disorder	4, 2.9
Hysteria	1, 0.7
Tic	3, 2.2
Sighing	3, 2.2
Speech delay	5, 3.7
Seizure	2, 1.5

One hundred five patients underwent medical treatment (77.2%), one of them underwent cardiac catheterization (0.7%), and 30 of them underwent surgery (22.1%). We followed the patients; in their last follow-up, 96 were stable, and 40 were missing the follow-up. There was no significant correlation between ADHD and congenital cardiac anomaly or malformation in our study population ($P > 0.05$). ADHD in CHD patients was less significant in any sex ($P > 0.05$). There was a significant correlation between prematurity and developmental delay ($P = 0.01$). The mean age was significantly different in patients with minor rather than those with major cardiac disorders ($P < 0.05$).

Discussion

Our study showed that ADHD was prevalent in CHD patients (31.6%) in the normal population. We also demonstrated that there was no significant correlation between ADHD

and special cardiac anomalies and is not sex-dependent. Moreover, there was a significant correlation between prematurity and developmental delay, and the mean age was significantly different in patients with minor rather than patients with major cardiac disorders. Although our study was not in a casualty design, prior investigations have suggested that CHD can be a risk factor for ADHD and autism. In a cross-sectional study, a significant proportion of children between the ages of 7 and 15 years old with CHD had symptoms of ADHD. Their results suggest that the prevalence of ADHD symptoms was significantly higher in this group with CHD (11.8%) when compared to the estimated prevalence in children (5%). In another study, the mean ADHD score in patients with CHD was the same in the general population with ADHD (22). Razzaghi et al. found that ADHD and autism were higher in children with CHD

than in the control group (23). Another study determined that the number of children with clinically significant ADHD scores among a patient with CHD was 3-4 times higher than that observed in the general population (24). Primary investigations mentioned that patients with CHD usually manifested unusual hemodynamic changes (25,26). This can lead to impaired cerebral blood flow (26) and abnormality in neurodevelopment (27-29), and immunological dysregulation (30,31), which has played an important role in the development of ADHD and autism (32,33). It was found in another study that the risk of ADHD and autism spectrum disorder was higher in children with developmental delays (34). Flanagan et al. reported that motor delays at the age of 6 months were associated with the development of autism at the age of 36 months (35). Moreover, Sharma et al. reported that young children with early developmental disorders were more likely to be subsequently diagnosed with ADHD (31). In another study, the first logistic regression model showed that children with CHD, compared with children without CHD, had ~1.6 to 2.7 times the odds of ADHD (36). According to previous studies, children with CHD are at higher risk of development and intelligence delay, with a higher incidence of gross and fine motor abnormalities and lower mean IQ scores than the age-matched controls (37,38). Marino et al. demonstrated that children with CHD were at increased risk of developmental disorders/disabilities and developmental delays (6). Therefore, it is important to identify the risk for ADHD and autism in a highly vulnerable subpopulation with two chronological risk factors, one present at birth and the other developing during early childhood. Indeed, the presence of one risk factor led to modest increases in risk for ADHD or autism. However, the hazard ratio for ADHD exponentially increased by 16.59 times (34). These results suggested that the existence of CHD can predispose the brain areas to dysfunction.

CHD in the neonatal period can cause neonates to suffer adverse perinatal outcomes, including perinatal infection, preterm delivery, or low birth weight (39). The potential role of these perinatal comorbid disorders that may have a negative impact on the development of ADHD (40) is still under debate. However, the current study did not show any correlation between comorbid perinatal disorders and ADHD. An increased incidence of ADHD in the

CHD patient population may be because of a higher prevalence of genetic syndromes in children undergoing heart surgery than in the normal population. Therefore, the risk for ADHD will, unfortunately, always be higher than in the normal population. However, an additional consideration is the potential impact of interrupting cerebral blood flow during neonatal aortic arch surgery with the subsequent loss of dopamine receptor activity (24). Reduced dopaminergic activity in the striatum has been documented in ADHD patients, indicating the possibility of a link between cardiac surgery, hypoxia, and attention-deficit/hyperactivity disorder (41).

Our study was a cohort study that enrolled CHD patients and followed them up. In this study, we missed follow-up in many patients. This study could be re-analyzed, and sub-group analysis can help to calculate hazard ratios of the Kaplan-Mayer test to achieve a more practical tool for predicting ADHD in CHD patients.

Conclusion

Attention deficit hyperactivity disorder has recently received more attention and is commonly seen in congenital heart disease. Cardiopulmonary signs and symptoms, abnormal electrocardiogram findings, and other associated neuropsychiatric disorders were significantly higher in these children. This study demonstrated that children born with congenital heart disease are at increased risk of suffering from attention deficit hyperactivity disorder. Further studies with a larger sample size and more formal attention deficit hyperactivity disorder evaluation can help evaluate the prevalence and risk factors in children with congenital heart disease.

A multidisciplinary approach consisting of a pediatrician, pediatric cardiologist, pediatric psychiatrist, and other related specialties is expected for timely and appropriate diagnosis and treatment.

Acknowledgments

The authors thank the Research and Technology Deputy of Mashhad University of Medical Sciences for financial support. The institutional review board approved this study at Mashhad University of Medical Sciences, Mashhad, Iran (IR.MUMS.MEDICAL.REC.1397.743). The authors declare any conflict of interest.

References

1. Zhao L, Chen L, Yang T, Wang T, Zhang S, Chen L, et al. Birth prevalence of congenital heart disease in China, 1980–2019: A systematic review and meta-analysis of 617 studies. *Eur J Epidemiol* 2020; 35(7): 631-42.
2. Griffin KJ, Elkin TD, Smith CJ. Academic outcomes in children with congenital heart disease. *Clin Pediatr* 2003; 42(5): 401-9.
3. Wray J, Sensky T. Congenital heart disease and cardiac surgery in childhood: Effects on cognitive function and academic ability. *Heart* 2001; 85(6): 687-91.
4. Rosenthal A, Castaneda AR. Growth and development after cardiovascular surgery in infants and children. *Prog Cardiovasc Dis* 1975; 18(1): 27-37.
5. Bass JL, Corwin M, Gozal D, Moore C, Nishida H, Parker S, et al. The effect of chronic or intermittent hypoxia on cognition in childhood: A review of the evidence. *Pediatrics* 2004; 114(3): 805-16.
6. Marino BS, Lipkin PH, Newburger JW, Peacock G, Gerdes M, Gaynor JW, et al. Neurodevelopmental outcomes in children with congenital heart disease: Evaluation and management: A scientific statement from the American Heart Association. *Circulation* 2012; 126(9): 1143-72.
7. Diamond A. Executive functions. *Ann Rev Psychol* 2013; 64: 135-68.
8. Blair C, Razza RP. Relating effortful control, executive function, and false belief understanding to emerging math and literacy ability in kindergarten. *Child Dev* 2007; 78(2): 647-63.
9. Diamond A, Barnett WS, Thomas J, Munro S. Preschool program improves cognitive control. *Science (New York, NY)* 2007; 318(5855): 1387.
10. Bellinger DC, Wypij D, Rivkin MJ, DeMaso DR, Robertson Jr RL, Dunbar-Masterson C, et al. Adolescents with d-transposition of the great arteries corrected with the arterial switch procedure: Neuropsychological assessment and structural brain imaging. *Circulation* 2011; 124(12): 1361-9.
11. Bellinger DC, Watson CG, Rivkin MJ, Robertson RL, Roberts AE, Stopp C, et al. Neuropsychological status and structural brain imaging in adolescents with single ventricle who underwent the Fontan procedure. *J Am Heart Assoc* 2015; 4(12): e002302.
12. Gaynor JW, Ittenbach RF, Gerdes M, Bernbaum J, Clancy RR, McDonald-McGinn DM, et al. Neurodevelopmental outcomes in preschool survivors of the Fontan procedure. *J Thorac Cardiovasc Surg* 2014; 147(4): 1276-83.
13. Cassidy AR, White MT, DeMaso DR, Newburger JW, Bellinger DC. Executive function in children and adolescents with critical cyanotic congenital heart disease. *J Int Neuropsychol Soc* 2015; 21(1): 34-49.
14. Calderon J, Jambaqué I, Bonnet D, Angeard N. Executive functions development in 5-to 7-year-old children with transposition of the great arteries: A longitudinal study. *Dev Neuropsychol* 2014; 39(5): 365-84.
15. Yamada DC, Porter AA, Conway JL, LeBlanc JC, Shea SE, Hancock-Friesen CL, et al. Early repair of congenital heart disease associated with increased rate of attention deficit hyperactivity disorder symptoms. *Can J Cardiol* 2013; 29(12): 1623-8.
16. Goldberg CS, Mussatto K, Licht D, Wernovsky G. Neurodevelopment and quality of life for children with hypoplastic left heart syndrome: current knowns and unknowns. *Cardiol Young* 2011; 21 Suppl 2(02): 88-92.
17. Neal AE, Stopp C, Wypij D, Bellinger DC, Dunbar-Masterson C, DeMaso DR, et al. Predictors of health-related quality of life in adolescents with tetralogy of Fallot. *J Pediatr* 2015; 166(1): 132-8.
18. DeMaso DR, Labella M, Taylor GA, Forbes PW, Stopp C, Bellinger DC, et al. Psychiatric disorders and function in adolescents with d-transposition of the great arteries. *J Pediatr* 2014; 165(4): 760-6.
19. Luyckx K, Rassart J, Goossens E, Apers S, Oris L, Moons P. Development and persistence of depressive symptoms in adolescents with CHD. *Cardiol Young* 2016; 26(6): 1115-22.
20. Matte B, Anselmi L, Salum GA, Kieling C, Gonçalves H, Menezes A, et al. ADHD in DSM-5: A field trial in a large, representative sample of 18- to 19-year-old adults. *Psychol Med* 2015; 45(2): 361-73.
21. Hansen E, Poole TA, Nguyen V, Lerner M, Wigal T, Shannon K, et al. Prevalence of ADHD symptoms in patients with congenital heart disease. *Pediatr Int* 2012; 54(6): 838-43.
22. Kovacs AH, Saidi AS, Kuhl EA, Sears SF, Silversides C, Harrison JL, et al. Depression and anxiety in adult congenital heart disease: predictors and prevalence. *Int J Cardiol* 2009; 137(2): 158-64.
23. Razzaghi H, Oster M, Reefhuis J. Long-term outcomes in children with congenital heart disease: National Health Interview Survey. *J Pediatr* 2015; 166(1): 119-24.

24. Shillingford AJ, Glanzman MM, Ittenbach RF, Clancy RR, Gaynor JW, Wernovsky G. Inattention, hyperactivity, and school performance in a population of school-age children with complex congenital heart disease. *Pediatrics* 2008; 121(4): e759-e67.
25. Rosenthal GL. Patterns of prenatal growth among infants with cardiovascular malformations: Possible fetal hemodynamic effects. *Am J Epidemiol* 1996; 143(5): 505-13.
26. Donofrio M, Bremer Y, Schieken R, Gennings C, Morton L, Eidem BW, et al. Autoregulation of cerebral blood flow in fetuses with congenital heart disease: the brain sparing effect. *Pediatr Cardiol* 2003; 24(5): 436-43.
27. Rollins CK, Watson CG, Asaro LA, Wypij D, Vajapeyam S, Bellinger DC, et al. White matter microstructure and cognition in adolescents with congenital heart disease. *J Pediatr* 2014; 165(5): 936-44.
28. Khalil A, Bennet S, Thilaganathan B, Paladini D, Griffiths P, Carvalho JS. Prevalence of prenatal brain abnormalities in fetuses with congenital heart disease: A systematic review. *Ultrasound Obstet Gynecol* 2016; 48(3): 296-307.
29. Donofrio MT, Massaro AN. Impact of congenital heart disease on brain development and neurodevelopmental outcome. *Int J Pediatr* 2010; 2010: 359390.
30. Madhok AB, Ojamaa K, Haridas V, Parnell VA, Pahwa S, Chowdhury D. Cytokine response in children undergoing surgery for congenital heart disease. *Pediatr Cardiol* 2006; 27(4): 408-13.
31. Sharma R, Bolger AP, Li W, Davlouros PA, Volk H-D, Poole-Wilson PA, et al. Elevated circulating levels of inflammatory cytokines and bacterial endotoxin in adults with congenital heart disease. *Am J Cardiol* 2003; 92(2): 188-93.
32. Oades RD, Dauvermann MR, Schimmelmann BG, Schwarz MJ, Myint A-M. Attention-deficit hyperactivity disorder (ADHD) and glial integrity: S100B, cytokines and kynurenine metabolism-effects of medication. *Behav Brain Funct* 2010; 6(1): 1-14.
33. Wei H, Zou H, Sheikh AM, Malik M, Dobkin C, Brown WT, et al. IL-6 is increased in the cerebellum of autistic brain and alters neural cell adhesion, migration and synaptic formation. *J Neuroinflammation* 2011; 8: 52.
34. Tsao P-C, Lee Y-S, Jeng M-J, Hsu J-W, Huang K-L, Tsai S-J, et al. Additive effect of congenital heart disease and early developmental disorders on attention-deficit/hyperactivity disorder and autism spectrum disorder: A nationwide population-based longitudinal study. *Eur Child Adolesc Psychiatry* 2017; 26(11): 1351-9.
35. Flanagan JE, Landa R, Bhat A, Bauman M. Head lag in infants at risk for autism: A preliminary study. *Am J Occup Ther* 2012; 66(5): 577-85.
36. Gonzalez VJ, Kimbro RT, Cutitta KE, Shabosky JC, Bilal MF, Penny DJ, et al. Mental health disorders in children with congenital heart disease. *Pediatrics* 2021; 147(2): e20201693.
37. Mari MA, Cascudo MM, Alchieri JC. Congenital heart disease and impacts on child development. *Braz J Cardiovasc Surg* 2016; 31(1): 31-7.
38. Lata K, Mishra D, Mehta V, Juneja M. Neurodevelopmental status of children aged 6–30 months with congenital heart disease. *Indian Pediatr* 2015; 52(11): 957-60.
39. Almeida LFG, Araujo Júnior E, Crott GC, Okido MM, Berezowski AT, Duarte G, et al. [Epidemiological risk factors and perinatal outcomes of congenital anomalies]. *Rev Bras Ginecol Obstet* 2016; 38(7): 348-55. (Portuguese)
40. Pringsheim T, Sandor P, Lang A, Shah P, O'Connor P. Prenatal and perinatal morbidity in children with Tourette syndrome and attention-deficit hyperactivity disorder. *J Dev Behav Pediatr* 2009; 30(2): 115-21.
41. Decker MJ, Rye DB. Neonatal intermittent hypoxia impairs dopamine signaling and executive functioning. *Sleep Breath* 2002; 6(4): 205-10.